



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Factors Influencing Adoption of Cattle Excreta Management Practices for Improved Elephant Grass (*Pennisetum purpureum*) Production by Smallholder Dairy Farmers

Fred Kabi
Makerere University, Uganda

F. B. Bareeba
Makerere University, Uganda

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/13-1/37>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Factors influencing adoption of cattle excreta management practices for improved elephant grass (*Pennisetum purpureum*) production by smallholder dairy farmers

F. Kabi ^{a*} and F.B. Bareeba ^a

^{a*} Department of Animal Science, Faculty of Agriculture, Makerere University, P.O. Box 7062 Kampala, Uganda, fredkabi@agric.mak.ac.ug

Key words: adoption, cattle excreta management, elephant grass production, smallholder dairy farmer

Introduction Livestock in East Africa have been reported to play a valuable role in sustainable agricultural systems (Bebe, 2003; Winrock International, 1992) where together with crops they make sustainable natural resource use possible (Lekasi et al., 2001). Haque et al. (1995) reported that animal excreta plays an important role in maintaining cropland productivity in Sub-Saharan Africa, and most smallholder farmers in Uganda will continue to rely on the exploitation of this natural resource for maintaining soil fertility. With the ever increasing human population and escalating demand for foods of animal origin, sustainable elephant grass production is of paramount importance among smallholder zero-grazing (intensive dairy) farmers in peri-urban areas of Uganda. However, in Uganda, elephant grass is becoming less productive both in dry matter (DM) yield and nutritive quality with successive harvesting. The decline in quality and quantity of elephant grass has been attributed to failure by the farmers to adopt sustainable livestock production practices (Katuromunda et al., 2001). Therefore, the objective of this study was to assess factors that influence adoption of cattle excreta management practices for improved elephant grass production among the intensive dairy farmers.

Materials and methods A diagnostic survey and participatory on-farm trials were conducted among intensive dairy farmers to examine factors that influence adoption of cattle excreta management practices. Management practices of burying cattle excreta in trenches, topical application and no applied excreta were compared. Influence of excreta management practices on quality and quantity of elephant grass were studied.

Results and discussion Although 98% of the sampled households were 5-7 people or more, 72% practiced zero-grazing on 2 acres or less. Therefore, 55% of the farmers kept only 2 crossbred dairy cows producing limited excreta for farm use, and 78% of the farmers produced less than 10 litres of milk from each cow. Burying cattle excreta resulted in higher ($P < 0.05$) dry matter (DM) yields of 22.4 t ha⁻¹ of elephant grass leaves compared to 15.1 t ha⁻¹ and 10.5 t ha⁻¹ from topical excreta application and control treatments, respectively. Although crude protein (CP g kg⁻¹) did not vary ($P > 0.05$) with the different cattle excreta management practices, DM yield of elephant grass, its leaf:stem ratio and ash were superior ($P < 0.05$) with treatments where the excreta was buried. While burying excreta improved quality and quantity of elephant grass, yields varied from farm to farm depending on the level of adoption of cattle excreta management practice. High cost, unfair division of labour among gender and lack of basic equipment such as spades and wheelbarrows to carry the excreta to elephant grass gardens affected the adoption of cattle excreta management practices. The major cost of production among zero-grazing farmers was feed, which accounted for 46.8% of total cost, followed by labour, veterinary services and equipment. It is concluded that high costs and unfair division of labour between gender as well as lack of basic equipment to transport excreta influenced the adoption of cattle excreta use by zero-grazing farmers.

Conclusion These results indicated that interventions aimed at improving elephant grass production by smallholder dairy farmers can only be successful if basic equipment that enhance recycling the excreta and gender roles to reduce the drudgery of labour are addressed.

References

- Bebe, B.O., 2003. Herd dynamics of smallholder dairy in Kenya highlands. *Ph.D. Thesis, Wageningen University, The Netherlands*, 155 pp.
- Haque, I., Powell, M.J., Ehui, K.S., 1995. Improved crop-livestock production strategies for sustainable soil management in tropical Africa. In: Lal, R., Steward, B.A. (eds.), *Advances in Soil Science and Soil Management. Experimental Basis for Sustainability and Environmental Quality*. Boca Raton: CRC Lewis Publishers, 293-345.
- Katuromunda, S., Sabiiti, E.N., Bareeba, F.B., 2001. Sustainable *Pennisetum purpureum* fodder production in the peri-urban smallholder crop-livestock production systems of Uganda. *African Crop Science Journal* 5, 559-564.